

ATI RN LEADERSHIP PROCTORED (3 LATEST VERSIONS) VERSION A, B AND C
LATEST | UPDATE|COMPREHENSIVE QUESTIONS AND VERIFIED
SOLUTIONS/CORRECT ANSWERS|GET 100% ACCURATE!!

A client with peptic ulcer disease has a new NG tube in place with orders for NG tube monitoring for the nurse. What is an expected finding? - ✓✓ANSWER✓✓>>scant blood may be seen for the first 12-24 hrs

A nurse has prepared a sterile field for assisting a provider with a chest tube insertion. Which of the following events should the nurse recognize as contaminating the field (Select all that apply)

- A. provider drops a sterile instrument onto the near side of the sterile field
 - B. nurse moistens a cotton ball with sterile normal saline and places it on sterile field
 - C. procedure is delayed 1hr because the provider receives an emergency call
 - D. nurse turns to speak to someone who enters through the door behind the nurse
 - E. clients hand brushes against the outer edge of the sterile field -
- ✓✓ANSWER✓✓>>B, C, D

A nurse has prepared a sterile field for assisting a provider with a chest tube insertion. Which of the following events should the nurse recognize as contaminating the field (Select all that apply)

- A. provider drops a sterile instrument onto the near side of the sterile field
 - B. nurse moistens a cotton ball with sterile normal saline and places it on sterile field
 - C. procedure is delayed 1hr because the provider receives an emergency call
 - D. nurse turns to speak to someone who enters through the door behind the nurse
 - E. clients hand brushes against the outer edge of the sterile field -
- ✓✓ANSWER✓✓>>B, C, D

A nurse has removed a sterile pack from its outside cover and place it on a clean work surface in preparation for an invasive procedure. Which of the following flaps should the nurse unfold first:

- A. closest to body
- B. right side
- C. left side
- D. farthest from body - ✓✓ANSWER✓✓>>D

A nurse has removed a sterile pack from its outside cover and place it on a clean work surface in preparation for an invasive procedure. Which of the following flaps should the nurse unfold first:

- A. closest to body
- B. right side
- C. left side
- D. farthest from body - ✓✓ANSWER✓✓>>D

A nurse is wearing sterile gloves in prep for performing a sterile procedure. Which of the following objects can the nurse touch without breaking sterile technique (Select all that apply)

- A. bottle containing sterile solution
- B. edge of sterile drape at the base of the field
- C. inner wrapping of an item on the sterile field
- D. irrigation syringe on the sterile field
- E. one gloved hand with the other gloved hand - ✓✓ANSWER✓✓>>C, D, E

A nurse is wearing sterile gloves in prep for performing a sterile procedure. Which of the following objects can the nurse touch without breaking sterile technique (Select all that apply)

- A. bottle containing sterile solution
- B. edge of sterile drape at the base of the field
- C. inner wrapping of an item on the sterile field
- D. irrigation syringe on the sterile field
- E. one gloved hand with the other gloved hand - ✓✓ANSWER✓✓>>C, D, E

A pt comes to the hospital with a TB induration of 10mm what would be your following nursing action? - ✓✓ANSWER✓✓>>obtain sputum cultures

A pt has a positive TB skin test what would your first action be? - ✓✓ANSWER✓✓>>put a mask on either the pt or yourself

A pt has a positive TB skin test what would your first action be? - ✓✓ANSWER✓✓>>put a mask on either the pt or yourself

ACE inhibitors side effects: - ✓✓ANSWER✓✓>>A: angioedema

- C: cough
- E: elevated K+

Acute hemolytic transfusion reaction s/s: - ✓✓ANSWER✓✓>>-chills

- fever
- low back pain
- tachycardia
- flushing
- hypotension

- chest tightening/pain
- tachypnea
- nausea
- anxiety
- hemoglobinuria

Anaphylactic transfusion reaction s/s: - ✓✓ANSWER✓✓>>-wheezing

- dyspnea
- chest tightness
- cyanosis
- hypotension

Appropriate oropharyngeal suctioning; select all: - ✓✓ANSWER✓✓>>-encourage pt to deep breath & cough prior

- monitor O2 sats
- store catheter in clean dry place for use
- perform care 3 times a day (Q 8hrs)

Appropriate oropharyngeal suctioning; select all: - ✓✓ANSWER✓✓>>-encourage pt to deep breath & cough prior

- monitor O2 sats
- store catheter in clean dry place for use
- perform care 3 times a day (Q 8hrs)

Asthma & dry powder inhaler teaching: - ✓✓ANSWER✓✓>>breath in forcefully

Automated external defibrillator steps: - ✓✓ANSWER✓✓>>1. turn it on & place pads on client

2. make sure no one is touching client
3. hit analyze
4. if shock is needed, AED will do it

AV block - ✓✓ANSWER✓✓>>AV block

AV block - ✓✓ANSWER✓✓>>AV block

Before an Endoscopy (EGD): - ✓✓ANSWER✓✓>>-NPO 6-8 hr

- remove dentures

Before an Endoscopy (EGD): - ✓✓ANSWER✓✓>>-NPO 6-8 hr

- remove dentures

Bilateral limb involvement occurs, making walking/balancing difficult. Which stage of Parkinson's? - ✓✓ANSWER✓✓>>Stage 2

brady - ✓✓ANSWER✓✓>>brady

brady - ✓✓ANSWER✓✓>>brady

Caring for a client following a stroke of the right side with left-sided hemipalegia: -

✓✓ANSWER✓✓>>-thicken liquids/foods

- high fowlers
- speech therapist for helping w/eating & speech
- monitor gag reflex/swallowing abilities
- have suction equipment available
- occupational therapy
- support left arm with pillows, slings, etc

Caring for a client following a stroke of the right side with left-sided hemipalegia: -

✓✓ANSWER✓✓>>-thicken liquids/foods

- high fowlers
- speech therapist for helping w/eating & speech
- monitor gag reflex/swallowing abilities
- have suction equipment available
- occupational therapy
- support left arm with pillows, slings, etc

Caring for a client who has pancreatitis: - ✓✓ANSWER✓✓>>-keep NPO

- NG tube to suction gastric contents
- low fat, high protein, high carb diet

Risk factors:

- high fat diet
- excessive alcohol consumption
- age

Caring for a client who has pancreatitis: - ✓✓ANSWER✓✓>>-keep NPO

- NG tube to suction gastric contents
- low fat, high protein, high carb diet

Risk factors:

- high fat diet
- excessive alcohol consumption
- age

Central venous access device care: - ✓✓ANSWER✓✓>>-remove tubing & solution bag every 24 hr

- DO NOT use line for other bolus IV solutions
- sterile dressing changes
- check cap glucose Q 4-6 hr