ATI COMPREHENSIVE EXIT EXAM 180 QUESTIONS AND ANSWERS VERIFIED AND WELL GRADED 2024, BEST ATI COMPREHENSIVE 2023 ATI RN PROCTORED COMPREHENSIVE PREDICTOR FORM A 180 QUESTIONS WITH ANSWERS (RATED A+)

> written by latestsolutions



Did you know a seller earn an average of \$250 per month selling their study notes on Docmerit



Scan the QR-code and learn how you can also turn your class notes, study guides into real cash today.

## Docmerit.com - The Best Study Notes

Uploaded by: latestsolutions on Docmerit. Distribution of this document is illegal

## 2023 ATI RN PROCTORED COMPREHENSIVE PREDICTOR FORMA 180 QUESTIONS WITH ANSWERS (RATED <u>A+)</u>

- 1. A nurse is caring for a client who is at 33 weeks of gestation following an amniocentesis. The nurse should monitor the client for which of the following complications?
  - a. Vomiting
  - **b.** Hypertension
  - c. Epigastric pain
  - d. Contractions
- 2. A nurse is providing teaching to an older adult client about methods to promote nighttime sleep. Which of the following instructions should the nurse include?
  - a. Stay in bed at least 1 hr if unable to fall asleep
  - b. Take a 1 hr nap during the day
  - c. Perform exercises prior to bedtime
  - d. Eat a light snack before bedtime
- 3. A nurse on a telemetry unit is caring for a client who becomes unconscious and whose monitor displays ventricular tachycardia. Which of the following actions should the nurse take first after determining the client does not have a palpable pulse?
  - a. Assess heart sounds
  - **b.** Defibrillate
  - c. Establish IV access
  - d. Administer epinephrine
- 4. A nurse is admitting a client who 1 week postpartum and reports excessive vaginal bleeding. The nurse does not speak the same language as the client. The client's partner and 10-year-old child are accompanying her. Which of the following actions should the nurse take to gather the client's admission data?
  - a. Have the client's child translate
  - **b.** Allow the client's partner to translate
  - c. Request a female interpreter through the facility
  - d. Ask a nursing student who speaks the same language as the client to translate
- 5. A nurse is caring for a client who is febrile (High fever). To reduce the client's fever, the nurse applies a cooling blanket. Which of the following findings indicates the client is having an adverse reaction to the cooling?
  - a. Flushing
  - b. Tachycardia
  - c. Restlessness
  - d. Shivering (Hypothermic)

6. A nurse is caring for a client who has deep-vein-thrombosis of the left lower extremity. Which of the following actions should the nurse take? (Exhibit)

|                                                       | Exhibit                                                                                                                                                  |                                            |                                         | 2         |
|-------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------|-----------------------------------------|-----------|
| tremity lower than the heart.                         | Tab1                                                                                                                                                     | Tab 2                                      | Tab3                                    |           |
| 4 hr.                                                 | Medicati<br>Record                                                                                                                                       | on Admini                                  | stration                                |           |
| aPTT 100 seconds<br>Platelets 140,000/mm <sup>3</sup> | Heparin IV infusion 1,000 units/hr<br>Warfarin 2 mg PO daily<br>Acetaminophen 650 mg PO every 4<br>hr PRN temperature greater than<br>38.6° C (101.5° F) |                                            |                                         |           |
| INR 2.0<br>10 비 요                                     | mo/aceta                                                                                                                                                 | one bitartri<br>minophen i<br>ter than 5 d | ate 5<br>500 mg PO Pi<br>on a scale fro | RN<br>m 0 |

- a. Position the client with the affected extremity lower than the heart
- b. Withhold heparin IV infusion PTT- 30-40 seconds; x2 if on heparin
- c. Administer acetaminophen
- d. Massage the affected extremity every 4 hr
- 7. A nurse is reviewing assessment data from several clients. For which of the following clients should the nurse recommend referral to a dietitian?
  - a. An older adult client who has a BMI of 24 (18.5-24.9)
  - b. A client who has a nonhealing leg ulcer (diet isn't good)
  - c. An older adult client who has presbyopia (age related far-sightness)
  - **d.** A client who has an albumin level of 3.7 g/dL (normal 3.4-5.4)
- 8. A nurse is providing discharge teaching to a client who has a chronic kidney disease and is receiving hemodialysis. Which of the following instructions should the nurse include in the teaching?
  - a. Eat 1 g/kg of protein per day
  - b. Take magnesium hydroxide for indigestion
  - c. Drink at least 3 L of fluid daily-
  - d. Consume foods high in potassium- restrict
- 9. A nurse is caring for a client who is receiving intermittent enteral tube feedings. Which of the following places the client at risk for aspiration?
  - **a.** Sitting in a high-Fowler's position during the feeding
  - b. A history of gastroesophageal reflux disease
  - c. Receiving a high osmolarity formula
  - d. A residual of 65 mL 1 hr postprandial?
- 10. ?A nurse is providing prenatal teaching to a client who is at 12 weeks of gestation. The nurse should tell the client that she will undergo which of the following screening tests at 16 weeks of gestation?
  - a. Chorionic villus sampling- as early as 8 weeks
  - **b.** Cervical cultures for chlamydia- 1<sup>st</sup> appointment.
  - c. Nonstress test -28 weeks
  - d. Maternal serum alpha-fetoprotein- 16 to 18 weeks